

Open Access

Recent Advanced Endoscopic Management of Endoscopic Retrograde Cholangiopancreatography Related Duodenal Perforations

Seon Mee Park

Department of Internal Medicine, Chungbuk National University College of Medicine, Cheongju, Korea

The management strategy for endoscopic retrograde cholangiopancreatography-related duodenal perforation can be determined based on the site and extent of injury, the patient's condition, and time to diagnosis. Most cases of perivaterian or bile duct perforation can be managed with a biliary stent or nasobiliary drainage. Duodenal wall perforations had been treated with immediate surgical repair. However, with the development of endoscopic devices and techniques, endoscopic closure has been reported to be a safe and effective treatment that uses through-the-scope clips, ligation band, fibrin glue, endoclips and endoloops, an over-the-scope clipping device, suturing devices, covering luminal stents, and open-pore film drainage. Endoscopic therapy could be instituted in selected patients in whom perforation was identified early or during the procedure. Early diagnosis, proper conservative management, and effective endoscopic closure are required for favorable outcomes of non-surgical management. If endoscopic treatment fails, or in the cases of clinical deterioration, prompt surgical management should be considered. **Clin Endosc 2016;49:376-382**

Key Words: Cholangiopancreatography, endoscopic retrograde; Perforation; Endoscopic closure

INTRODUCTION

Endoscopic retrograde cholangiopancreatography (ERCP)-related duodenal perforation develops in 0.4% to 0.6% of patients, with serious adverse effects on morbidity and mortality.^{1,2} With the improvement of the endoscopic and surgical treatments of duodenal perforation, the outcomes have improved.^{3,4} The clinical outcomes depend on the timing of the diagnosis and the efficacy of the treatment. Treatment strategy depends on the type of perforation such as perivaterian or ductal defects managed with nonsurgical management, or duodenal wall perforation treated with surgical repair.^{4,5}

However, with the development of endoscopic devices and techniques, the success rate of endoscopic closure in duodenal wall perforation was reported to be as high as 88% in a recent meta-analysis.⁶ Endoscopic management is recommended for iatrogenic gastrointestinal perforation in selected patients with low level of evidence.² This review aimed to introduce advanced endoscopic management for ERCP-related duodenal perforation and to define its optimal management strategies.

CLASSIFICATION AND RISK FACTORS

Perforations related with ERCP were classified depending on their locations as follows: duodenal wall by the scope itself (type I), perivaterian by endoscopic sphincterotomy (EST) or endoscopic transpapillary balloon dilation (EPBD; type II), bile duct by a guidewire or basket (type III), and retroperitoneal air alone by compressed air leakage (type IV) (Fig. 1).⁷ This classification is widely used because it is appropriately related with clinical severity, perforation mechanism, anatom-

Received: June 26, 2016 Revised: July 18, 2016

Accepted: July 18, 2016

Correspondence: Seon Mee Park

Department of Internal Medicine, Chungbuk National University College of Medicine, 1 Chungdae-ro, Seowon-gu, Cheongju 28644, Korea
 Tel: +82-43-269-6019, Fax: +82-43-273-3252, E-mail: smpark@chungbuk.ac.kr

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ical location, and the need of surgical management.⁷ The main causes of perforation are EST in 41%, the endoscope itself in 26%, the guidewire in 15%, dilation of a stenotic segment in 3%, ERCP devices in 4%, and stent insertion and migration in 2%.⁴

The common risk factors related to duodenal perforation are old age and long procedure time.⁵ Specific risk factors are altered surgical anatomy, including Billroth II gastrectomy and Roux-en-Y bypass surgery, in duodenal wall perforations, and biliary stricture dilation, sphincter of Oddi dysfunction, and endoscopic papillectomy in perivaterian or ductal perforations.^{5,8,9} Whether diverticulum in perivaterian or needle-type papillotome is a risk factor remains controversial.^{2,10} EST and EPBD showed the same risk of perforation in a meta-analysis.¹¹




Fig. 1. Classification of endoscopic retrograde cholangiopancreatography-related duodenal perforation. Type I, lateral or medial wall duodenal perforation; type II, periampullary perforation; type III, bile duct injuries; type IV, retroperitoneal air alone.

RECOGNITION OF DUODENAL PERFORATION

Prompt recognition of duodenal perforation during a procedure is mandatory for favorable prognosis. However, many studies reported that 30% to 44.6% of perforations were found after the procedure.⁴ Duodenal wall perforations can be detected by direct visualization on endoscopy, and perivaterian or ductal perforations are usually recognized on fluoroscopy.¹² The use of carbon dioxide for inflation may minimize extraluminal air; therefore, free air or retroperitoneal air is difficult to detect during the procedure.⁴ Perforations should be suspected in the presence of abdominal pain, fever, and leukocytosis, even if a perforation was not detected at routine chest and abdominal radiography during or after the procedure. Clinical signs sometimes develop late, with peritoneal irritation signs and leukocytosis appearing at 3 to 4 and 12 hours after the procedure, respectively.⁴ In addition, perforation is misdiagnosed as post-ERCP pancreatitis or abdominal distension with inflated air. Forty-three percent of patients with perforation revealed a concurrent post-ERCP pancreatitis.¹³ Therefore, when perforation is suspected, a contrast-enhanced abdominal computed tomography (CT) scan should be performed, which is the most useful tool to detect perforation and provides useful information about pancreatitis and fluid collection (Fig. 2).¹⁴

When a duodenal perforation is detected endoscopically, a comprehensive examination and clear report should be written about its size and location with a picture, the endoscopic treatment, whether carbon dioxide or air was used for insufflation, and the standard report information. Because air inflation itself during endoscopy may not worsen the clinical outcome, comprehensive endoscopic examination is essential for proper treatment.²




Fig. 2. Computed tomography (CT) findings indicating a duodenal perforation. (A) An abdominal CT image showing massive fluid and air collection at the retroperitoneal space (type II perforation). (B) An abdominal CT image showing massive air leakage without fluid accumulation in the retroperitoneal space (type IV perforation).




Fig. 3. Endoscopic closure of a duodenal wall perforation by using ligation band and endoclips. (A) A large duodenal perforation on the lateral wall. (B) Endoscopic closure with band ligation and endoclips. (C) Endoscopic band ligation.




Fig. 4. Endoscopic closure of an anastomotic perforation by using an over-the-scope clipping device (OTSC). (A) OTSCs. (B) Jejunum perforation caused by Billroth II anastomotic leak. (C) Using a twin grasping forceps, the edges of the perforation wall are pulled back into the lumen of the cap attached to the endoscope. (D) The perforation site is successfully sealed with an OTSC.

ENDOSCOPIC MANAGEMENT

Duodenal wall perforation

Duodenal wall perforations require early closure because

large amounts of intestinal contents leak into the peritoneum. Endoscopic techniques are determined based on the type and size of perforation and the endoscopist expertise available at the center.^{15,16} The most important step before the closing

procedure is thorough aspiration of the duodenal contents through the perforated hole.

A duodenal perforation of <10 mm in diameter can be closed easily with endoclips¹⁷ by using a single-channel endoscope or lateral scope without changing the scope. Through-the-scope (TTS) clips are widely used for closing luminal perforations,¹⁸ achieving technical success in 90% of gut perforations.⁶ However, a large perforation (>20 mm) located at angulation is difficult to close by using TTS clips alone. When luminal closure with TTS clips fail, alternative methods using fibrin glue,¹⁹ endoclips and endoloops,²⁰ purse-string suture,²¹ or band ligation²² are applied.^{15,16} Fibrin glue spray through a standard ERCP catheter between the clips could obliterate the perforation site and adhere multiple clips to each other.¹⁹ Multiple endoclips and single or double endoloops are also effective for endoscopic closure of large duodenal perforations.^{20,21} Another alternative method for duodenal closure is endoscopic band ligation (EBL).²² EBL could be easily performed for perforations 1 to 2 cm in diameter regardless of the site or angle when a perforation is detected during the procedure.²³ Gastroscopy attached with single- or multiple-band ligators in a transparent cap can approach the edge of the perforation. The perforated wall, along with peritoneal fat, is sucked into

the ligator cup. One or two bands are subsequently released, resulting in closure of the perforation. Sucking together the duodenal wall and nearby omentum is helpful for complete closure of the perforation by making an omental patch endoscopically.²⁴ When a residual defect is suspected, additional endoclips can be applied at the edges of the perforation (Fig. 3).²⁵ Duodenal wall perforation can be easily ligated because of a thin wall structure.²⁶ Endoscopic devices such as endoclip, endoloop, ligation band, and fibrin glue are usually available in an endoscopy unit and can be applied as soon as perforation occurs.

Specialized devices such as over-the-scope clipping device (OTSC),²⁷ endoscopic suturing (OverStitch; Apollo Endosurgery, Austin, TX, USA),²⁸ and open-pore film drainage²⁹ are reported as useful and effective for large luminal perforation.

OTSC revealed an 88% technical success rate.⁶ OTSCs can close defects by up to 20 mm. Detailed OTSC techniques are as follows: OTSC in an endoscopic cap is placed over the mucosal defect. A twin grasper captures opposite sides of an open defect and apposes the tissue by using left- and right-sided graspers. The tissue in the graspers is pulled into the cap device, and then a clip is applied (Fig. 4). Compared with the TTS clip, the OTSC has higher compression force and enables




Fig. 5. Vacuum-assisted closure of the esophageal defect due to an anastomotic leak. (A) Device made of porous sponge and attached to a nasogastric tube. (B) Positioning of sponge device at the esophageal defect. (C) After 3 days of therapy, the sponge device is filled with extraluminal fluid and gastrointestinal secretions. (D) Esophageal wall perforation before treatment. (E) Improved luminal defect with the sponge changed once after 3 days. (F) Closed wound with the sponge changed twice every 3 days.

capture of a larger tissue volume.³⁰ However, OTSC is not widely available and requires training before application in human patients.³¹

Endoscopic suturing devices such as Eagle Claw (Olympus, Tokyo, Japan) and Purse String Suturing Device (LSI Solutions, Victor, NY, USA), which are attached to the end of a therapeutic double channel endoscope, allow full-thickness tissue apposition.³² The device can be used multiple times without the need to remove the scope from the patient. However, endoscopic suturing devices are not widely available and are difficult to operate within the narrow duodenal lumen. They have been investigated in animals but have not been reported for clinical closure of duodenal perforations.³³

Another new technique is vacuum therapy with open-pore film drainage, which is used for duodenal perforations that occur during ERCP.²⁹ The procedure consists of placing a drainage tube, which is wrapped with an open pore film, in the duodenal wall defect and application of vacuum for evacuation and collapse of the stomach and duodenum. Gastric, biliary, and pancreatic liquids are drained in an intraluminal direction continuously, and the transmural defect is closed simultaneously; thereby, preventing further extraluminal contamination (Fig. 5).

Perivaterian or ductal perforation

Most cases of periampullary or ductal perforation are man-



Fig. 6. Abdominal computed tomography scan showing air collection in the periductal space (arrow). To block further bile and pancreatic juice leakage, a self-expandable stent (arrowhead) is inserted at the distal common bile duct.

aged with biliary diversion and endoscopic closure to prevent retroperitoneal fluid leakage.¹³ Biliary diversion is achieved with biliary stent or nasobiliary drainage. A fully covered self-expandable metal stent (FCSEMS) can be placed to drain the bile duct and occlude the leak site in perivaterian perforations (Fig. 6).³⁴ A recent study reported that the benefits of FCSEMS in the perivaterian perforation area are painless course, lower leukocyte counts, and short hospital stay compared with conservative treatment alone.³⁵ The efficacy of TTS clips in closing periampullary perforations is unknown. However, it is technically difficult and has a risk of clipping the ampulla.^{36,37}

Ductal perforations are usually small in size and improve with conservative management.³ Perforation of the distal bile duct often can be managed by placement of plastic stents and, more recently, by FCSEMS.¹³ If endoscopy fails to treat the perforation, percutaneous biliary drainage above the level of the perforation is an alternative treatment.²

Retroperitoneal air alone (type IV) is developed by compressed air leakage.^{38,39} It does not need treatment. Pneumoperitoneum can be present in up to 29% of asymptomatic patients who undergo uncomplicated procedures.⁴⁰

CONSERVATIVE MANAGEMENT

When a perforation occurs during an ERCP procedure, ERCP should be switched to carbon dioxide insufflations and decompression of tension pneumoperitoneum or tension pneumothorax before performing endoscopic procedures.¹⁵ Switching from air to carbon dioxide for insufflation in iatrogenic perforations may prevent tension-pneumothorax, tension-pneumomediastinum, or tension-pneumopericardium, and abdominal compartment syndrome. Tension pneumoperitoneum should be decompressed immediately with an 18- or 20-gauge needle near the umbilicus, with the patient in the supine position.²

Conservative managements are nil-by-mouth and intravenous administration of proton pump inhibitors, broad spectrum antibiotics, fluids, and on-demand pain medication. For the diversion of luminal content, nasogastric or nasoduodenal aspiration is used in all patients and total parenteral nutrition is needed. Cross-sectional imaging should be performed during follow-up, and if a liquid collection is disclosed, percutaneous drainage should be considered. Oral intake can be resumed after the fourth day if no further leakage is identified.²

THERAPEUTIC ALGORITHMS FOR ERCP-RELATED DUODENAL PERFORATIONS

When a duodenal wall perforation is detected during the procedure or earlier (<12 hours), endoscopic/conservative management could be performed.² When duodenal perforation is diagnosed later (>12 hours), endoscopic management could be performed in selected patients who are in good condition without extravasation of contrast medium or persistent large fluid collection on CT.² Perivaterian or ductal perforation could be started with nonsurgical management.¹⁶ During conservative management, surgery should be considered in the conditions of failed endoscopic closure, serious leakage of contrast material, development of sepsis during conservative management, peritonitis, undrained fluid retention in the peritoneum, and worsening patient condition.⁴¹ As salvage operation is related with worse outcomes than those in cases treated initially with operative repair, patients who require surgical repair at the time of diagnosis should be identified.¹

PROGNOSIS

The overall mortality of ERCP-related duodenal perforation is reported to be 7.8% to 9.9%.^{1,2} The prognosis of ERCP-related duodenal perforation is determined based on the type, interval between perforation and recognition, and early luminal closure to reduce serous fluid leak. The amount of extraluminal air is not proportional to the size of the perforation and clinical severity.⁴² Non-operative treatment was successful in 79% to 93% of patients with periampullary injuries and in all the patients with ductal perforation.^{3,32} Endoscopic closure for duodenal wall perforation was applied and succeeded clinically in only 22% of cases. However, the clinical success rate after successful endoscopic closure was 94%.² Patients treated with surgery showed poor prognosis with delayed diagnosis, salvage operation, repeated operation, or old age.¹

CONCLUSIONS

The incidence of ERCP-related duodenal perforation has decreased, and the outcomes of nonsurgical management have improved.^{1,4} This progression is achieved by early diagnosis, proper conservative management, and immediate endoscopic closure using various endoscopic devices. Endoscopic therapy could be instituted in all patients with intraprocedurally identified perforations.^{2,6} However, the mortality in ERCP-related perforations still remain at 7.8% and salvage operation was associated with poor prognosis.^{1,4} To reduce

morbidities and mortalities in ERCP-related perforations, evaluation of risk factors, early detection of perforation during ERCP procedures, and training in endoscopic closure techniques are mandatory. During nonsurgical management, a multidisciplinary approach by endoscopists, radiologists, and surgeons is recommended to determine when and who should receive operation.

Conflicts of Interest

The author has no financial conflicts of interest.

REFERENCES

- Machado NO. Management of duodenal perforation post-endoscopic retrograde cholangiopancreatography: when and whom to operate and what factors determine the outcome? A review article. *JOP* 2012;13:18-25.
- Paspatis GA, Dumonceau JM, Barthet M, et al. Diagnosis and management of iatrogenic endoscopic perforations: European Society of Gastrointestinal Endoscopy (ESGE) Position Statement. *Endoscopy* 2014;46:693-711.
- Andriulli A, Loperfido S, Napolitano G, et al. Incidence rates of post-ERCP complications: a systematic survey of prospective studies. *Am J Gastroenterol* 2007;102:1781-1788.
- Veidakis A, Fragulidis G, Polydorou A. Endoscopic retrograde cholangiopancreatography-related perforations: diagnosis and management. *World J Gastrointest Endosc* 2015;7:1135-1141.
- Enns R, Eloubeidi MA, Mergener K, et al. ERCP-related perforations: risk factors and management. *Endoscopy* 2002;34:293-298.
- Verlaan T, Voermans RP, van Berge Henegouwen MI, Bemelman WA, Fockens P. Endoscopic closure of acute perforations of the GI tract: a systematic review of the literature. *Gastrointest Endosc* 2015;82:618-628.
- Stapfer M, Selby RR, Stain SC, et al. Management of duodenal perforation after endoscopic retrograde cholangiopancreatography and sphincterotomy. *Ann Surg* 2000;232:191-198.
- Freeman ML, Nelson DB, Sherman S, et al. Complications of endoscopic biliary sphincterotomy. *N Engl J Med* 1996;335:909-918.
- Wu HM, Dixon E, May GR, Sutherland FR. Management of perforation after endoscopic retrograde cholangiopancreatography (ERCP): a population-based review. *HPB (Oxford)* 2006;8:393-399.
- Navaneethan U, Konjeti R, Venkatesh PG, Sanaka MR, Parsi MA. Early precut sphincterotomy and the risk of endoscopic retrograde cholangiopancreatography related complications: an updated meta-analysis. *World J Gastrointest Endosc* 2014;6:200-208.
- Zhao HC, He L, Zhou DC, Geng XP, Pan FM. Meta-analysis comparison of endoscopic papillary balloon dilatation and endoscopic sphincterotomy. *World J Gastroenterol* 2013;19:3883-3891.
- Motomura Y, Akahoshi K, Gibo J, et al. Immediate detection of endoscopic retrograde cholangiopancreatography-related periampullary perforation: fluoroscopy or endoscopy? *World J Gastroenterol* 2014;20:15797-15804.
- Kumbhari V, Sinha A, Reddy A, et al. Algorithm for the management of ERCP-related perforations. *Gastrointest Endosc* 2016;83:934-943.
- de Vries JH, Duijm LE, Dekker W, Guit GL, Ferwerda J, Scholten ET. CT before and after ERCP: detection of pancreatic pseudotumor, asymptomatic retroperitoneal perforation, and duodenal diverticulum. *Gastrointest Endosc* 1997;45:231-235.
- Kwon CI, Song SH, Hahm KB, Ko KH. Unusual complications related to endoscopic retrograde cholangiopancreatography and its endoscopic treatment. *Clin Endosc* 2013;46:251-259.
- Lee TH, Han JH, Park SH. Endoscopic treatments of endoscopic retro-

- grade cholangiopancreatography-related duodenal perforations. *Clin Endosc* 2013;46:522-528.
17. Lee TH, Bang BW, Jeong JL, et al. Primary endoscopic approximation suture under cap-assisted endoscopy of an ERCP-induced duodenal perforation. *World J Gastroenterol* 2010;16:2305-2310.
 18. Mangiavillano B, Viaggi P, Masci E. Endoscopic closure of acute iatrogenic perforations during diagnostic and therapeutic endoscopy in the gastrointestinal tract using metallic clips: a literature review. *J Dig Dis* 2010;11:12-18.
 19. Yang HY, Chen JH. Endoscopic fibrin sealant closure of duodenal perforation after endoscopic retrograde cholangiopancreatography. *World J Gastroenterol* 2015;21:12976-12980.
 20. Nakagawa Y, Nagai T, Soma W, et al. Endoscopic closure of a large ERCP-related lateral duodenal perforation by using endoloops and endoclips. *Gastrointest Endosc* 2010;72:216-217.
 21. Li Q, Ji J, Wang F, et al. ERCP-induced duodenal perforation successfully treated with endoscopic purse-string suture: a case report. *Oncotarget* 2015;6:17847-17850.
 22. Li Y, Han Z, Zhang W, Wang X, et al. Successful closure of lateral duodenal perforation by endoscopic band ligation after endoscopic clipping failure. *Am J Gastroenterol* 2014;109:293-295.
 23. Han JH, Lee TH, Jung Y, et al. Rescue endoscopic band ligation of iatrogenic gastric perforations following failed endoclip closure. *World J Gastroenterol* 2013;19:955-959.
 24. Angsuwatcharakon P, Thienchanachaiya P, Pantongrag-Brown L, Reknimitr R. Endoscopic band ligation to create an omental patch for closure of a colonic perforation. *Endoscopy* 2012;44 Suppl 2 UCTN:E90-E91.
 25. Fan CS, Soon MS. Repair of a polypectomy-induced duodenal perforation with a combination of hemoclip and band ligation. *Gastrointest Endosc* 2007;66:203-205.
 26. Ji JS, Cho YS. Endoscopic band ligation: beyond prevention and management of gastroesophageal varices. *World J Gastroenterol* 2013;19:4271-4276.
 27. Buffoli F, Grassia R, Iiritano E, Bianchi G, Dizioli P, Staiano T. Endoscopic "retroperitoneal fatpexy" of a large ERCP-related jejunal perforation by using a new over-the-scope clip device in Billroth II anatomy (with video). *Gastrointest Endosc* 2012;75:1115-1117.
 28. Kumta NA, Boumitri C, Kahaleh M. New devices and techniques for handling adverse events: claw, suture, or cover? *Gastrointest Endosc Clin N Am* 2015;25:159-168.
 29. Loske G, Rucktäschel F, Schorsch T, van Ackeren V, Stark B, Müller CT. Successful endoscopic vacuum therapy with new open-pore film drainage in a case of iatrogenic duodenal perforation during ERCP. *Endoscopy* 2015;47(Suppl 1):E577-E578.
 30. Donatelli G, Dumont JL, Vergeau BM, et al. Colic and gastric over-the-scope clip (Ovesco) for the treatment of a large duodenal perforation during endoscopic retrograde cholangiopancreatography. *Therap Adv Gastroenterol* 2014;7:282-284.
 31. Brodie M, Gupta N, Jonnalagadda S. Failed attempt at duodenal perforation closure with over-the-scope clip. *Gastrointest Endosc* 2015;81:1271-1272.
 32. Kumar N, Thompson CC. A novel method for endoscopic perforation management by using abdominal exploration and full-thickness sutured closure. *Gastrointest Endosc* 2014;80:156-161.
 33. Winder JS, Pauli EM. Comprehensive management of full-thickness luminal defects: the next frontier of gastrointestinal endoscopy. *World J Gastrointest Endosc* 2015;7:758-768.
 34. Cho KB. The management of endoscopic retrograde cholangiopancreatography-related duodenal perforation. *Clin Endosc* 2014;47:341-345.
 35. Odemis B, Oztas E, Kuzu UB, et al. Can a fully covered self-expandable metallic stent be used temporarily for the management of duodenal retroperitoneal perforation during ERCP as a part of conservative therapy? *Surg Laparosc Endosc Percutan Tech* 2016;26:e9-e17.
 36. Baron TH, Gostout CJ, Herman L. Hemoclip repair of a sphincterotomy-induced duodenal perforation. *Gastrointest Endosc* 2000;52:566-568.
 37. Katsinelos P, Paroutoglou G, Papaziosos B, Beltsis A, Dimiropoulos S, Atmatzidis K. Treatment of a duodenal perforation secondary to an endoscopic sphincterotomy with clips. *World J Gastroenterol* 2005;11:6232-6234.
 38. Singh V, Singh G, Verma GR, Yadav TD, Gupta V. Pseudobowel perforation following endoscopic retrograde cholangiopancreatography. *Dig Dis Sci* 2013;58:1781-1783.
 39. Jang JS, Lee S, Lee HS, et al. Efficacy and safety of endoscopic papillary balloon dilation using cap-fitted forward-viewing endoscope in patients who underwent Billroth II gastrectomy. *Clin Endosc* 2015;48:421-427.
 40. Genzlinger JL, McPhee MS, Fisher JK, Jacob KM, Helzberg JH. Significance of retroperitoneal air after endoscopic retrograde cholangiopancreatography with sphincterotomy. *Am J Gastroenterol* 1999;94:1267-1270.
 41. Alfieri S, Rosa F, Cina C, et al. Management of duodeno-pancreato-biliary perforations after ERCP: outcomes from an Italian tertiary referral center. *Surg Endosc* 2013;27:2005-2012.
 42. Baron TH, Wong Kee Song LM, Zielinski MD, Emura F, Fotoohi M, Kozarek RA. A comprehensive approach to the management of acute endoscopic perforations (with videos). *Gastrointest Endosc* 2012;76:838-859.